

Original Article

Smart Agriculture Using UAV Imagery and Deep Learning

PROF. DAVID R. MONTGOMERY

Department of Earth and Space Sciences, University of Washington, USA.

ABSTRACT: *Climate change, population growth, and resource scarcity present unprecedented challenges to global food security, demanding a paradigm shift toward precision agriculture. Labor-intensive, low temporal resolution, and atmospheric interference represented drawbacks associated with traditional crop monitoring methods involving manual field surveys or satellite remote sensing. This study presents a novel hybrid smart agriculture framework using Unmanned Aerial Vehicle (UAV) images and deep learning architectures to achieve high-resolution crop health monitoring and fully automated yield prediction. Multi-season datasets of high-resolution multispectral and digital aerial imagery were collected over an extensive area comprising a variety of experimental plots. The imagery was processed using a hybrid deep learning architecture that comprised a modified Convolutional Neural Network (CNN) to extract spatial features and incorporate them into a Long Short-Term Memory network for temporal sequence analysis. The framework combines multispectral vegetation indices with raw spatial information in two tasks: classifying crop health, localized nutrient deficiencies, and predicting final crop yields. Using experimental data as the illustrative showcase, we validate that a deep learning model achieves crop stress classification accuracy of 96.4% and yield prediction with $R^2 = 0.89$, outperforming traditional machine-learning models by a large margin on two axes (classification/regression). We develop a scalable and automated pipeline to enhance the translation of high resolution remote sensing into actionable agronomic insights, bridging disciplinary divides between computational science, plant biology, and agriculture such that it provides valuable tools for optimizing resource allocation in sustainable farms.*

KEYWORDS: Precision Agriculture, Unmanned Aerial Vehicles, Deep Learning, Convolutional Neural Networks, Remote Sensing, Yield Prediction.

1. INTRODUCTION

Today's agricultural industry is at a crossroads, caught between the dual pressures of feeding a burgeoning human population while reducing environmental impacts. Traditional uniform farming practices, spreading major quantities of water and fertilizers uniformly on massive areas coated with completely different crops, are already turning increasingly unsustainable. These practices result not just in inefficient resource utilization, but also greater production costs and grave environmental consequences like groundwater contamination as well as soil degradation. As a result, precision agriculture has become an important field as it intends to observe and evaluate inter- and intra-field variability in crops, followed by applying the necessary measures. Precision agriculture specifically provides site-specific input optimization, creating a potential mechanism for improving the sustainability and productivity of food systems by managing variability over space or time.

Traditionally, field monitoring consisted of visual inspections, which are also more subjective, difficult to implement, and impractical at scale. With the introduction of satellite remote sensing, these macro-level observations allowed periodic data to be captured over large areas. In satellite data, often single pixels will be made up of multiple land-cover types or crop conditions and thus have spatial resolution shortfalls for precision farming applications. This is compounded by the fact that satellites have fixed orbital schedules which sometimes leave certain vital growth phases of crops hidden under cloud cover, meaning farmers miss crucial data in their important decision-making times.

Uncrewed Aerial Vehicles (UAVs), or drones, provide a methodology for agricultural remote sensing that bridges the gap between ground and satellite observations. UAVs generate extremely high-spatial-resolution imagery features can be resolved to the centimeter range due in part to their low altitude of operation. The fine granularity enables the identification of single plants, early weed infestations, and small-scale variations in canopy architecture. Further, the operational versatility of UAVs allows users to time flights on-demand with pivotal phenological stages or react immediately following extreme weather while completely circumventing cloud cover needs. Under multispectral, hyperspectral, high-resolution digital sensors, UAVs have richly spectral data corresponding directly with biophysical or biochemical characteristics of crops.

While UAVs provide abundant spatial data, translating unprocessed imagery into agronomic insights is a major computational bottleneck. Traditional approaches to image processing depend on a substantial amount of hand-crafted features and empirically derived vegetation indices that usually do not accurately identify non-linear dynamics in complex agricultural ecosystems (Dandois et al. 2015). This limitation has led to the incorporation of computer vision and deep learning into precision agriculture. Deep learning algorithms are known for their power to automatically discover hierarchical feature representations from raw spatial data, such as Convolutional Neural Networks (CNNs). With several levels of computations stacked atop each other, these

networks can learn complex patterns along with textures and structural irregularities that cannot be picked up through traditional statistical models. Moreover, through the integration with recurrent architectures such as Long Short-Term Memory (LSTM) networks, deep learning can capture a complete season trajectory of crop growth over time.

The research tackles critical gaps in present smart farming workflows, proposing a holistic integrated all-in-one method combining multi-source UAV imagery with state-of-the-art deep learning models to concurrently map each crop stress classification and yield prediction. Whereas previous studies frequently consider weed detection, disease classification, and yield estimation as separate problems with disjointed solutions leveraging spatial, spectral, or temporal data streams independently from each other, we propose a system that processes these three focal areas holistically to output valuable decisions. Your holistic approach reflects the cumulative impacts of early-season stressors on final harvest metrics.

This study aims to:

- To appraise an automated end-to-end processing pipeline for multi-sensor UAV-retrieved imagery that a) radiometrically calibrates and spatially aligns geometric models between repetitive flights
- To develop a hybrid deep learning framework that combines the spatial-spectral characteristics obtained through convolutional layers and temporal dynamics captured using recurrent neural networks.
- Study the performance of the proposed framework to classify multi-class crop stresses (in this case, nutrient deficiency, water stress, and disease) under variable field conditions.
- It monitored and validated the predictive accuracy of the deep learning model concerning final crop yield from real harvester-collected data (colleagues have already published a baseline comparison against regular machine & human-supported methods).

This study makes a substantial contribution to the digital agronomy field by achieving those objectives. A multi-scale feature fusion method that preserves spatial information of early local crop anomalies while abstracting deep layers is proposed. We further offer extensive empirical data, recorded under various environmental regimes, to demonstrate that fusing multispectral indices with deep structural features enhances generalization capabilities compared to models using RGB input or only specific spectral bands. In the end, this is a practical tech roadmap away from reactionary farm practices to data-driven precision agriculture.

2. LITERATURE REVIEW

Remote sensing in agriculture has undergone an evolution defined by a search for increasingly higher spatial, spectral, and temporal resolution. Satellite platforms (Landsat and MODIS) dominated the early research landscape, which set most regional cropping monitoring systems and macro yield forecasting. Yet the early literature frequently emphasizes limitations in terms of site-specific management, as atmospheric attenuation and large pixel sizes smear spatial variability within field limits. Commercial constellations such as PlanetScope / Sentinel-2 provided enhanced temporal revisit periods and spatial resolutions, but the inherent vulnerability to cloud cover remains a lingering bottleneck during peak growing seasons in tropical/temperate regions.

UAV platforms emerged as a new option and opened up numerous opportunities for high-resolution characterization of canopy attributes around the early 2010s. Initial adoptions were focused on general computation of traditional Vegetation Indices (VIs), such as the NDVI and normalized difference red edge (NDRE) computed from low-cost multispectral sensors. They correlated these indices with chlorophyll, LAI, and biomass successfully. Even though these empirical regression models were demonstrated to work well under extremely tightly controlled experimental conditions, it was only when out-of-the-lab validation studies began being performed that real shortfalls became apparent. Early season growth estimations using empirical VIs are severely affected by soil background reflectance effects and typically saturate at high canopy densities, causing large errors in field health assessments late in the growing season as well as yield predictions.

However, since simple VIs suffer from saturation and linearity limitations, researchers turned to shallow machine learning algorithms. Various algorithms (Support Vector Machines SVM, Random Forests RF and Artificial Neural Networks ANN) started being the bread-and-butter of working with UAV-derived datasets. As an example, Random Forest classifiers were used to map weeds or for crop disease classification because of their ability to account for multi-collinear spectral bands in a non-overfitting way, among others. But these superficial machine learning tasks require substantial human feature engineering. Using this method, agronomists and data scientists had to hand-craft texture features, such as by applying the Gray-Level Co-occurrence Matrix (GLCM) or manual segmentation 1 leaf per row → biased human variability, which do not generalize well across species/location.

Naturally, the success of deep learning in computer vision made its way towards precision agriculture as well. Manual feature extraction was replaced by Convolutional Neural Networks (CNNs) that learned the spatial hierarchies of features automatically. In the early years of deep learning, we primarily saw applications in agriculture related to object detection (counting fruits or identifying specific weed plants within row crops using an architecture such as YOLO(You Only Look Once) or Faster-RCNN).

As computational capabilities expanded, researchers began applying pixel-level semantic segmentation architectures, such as U-Net and SegNet, to UAV orthomosaics to map the exact spatial extent of crop diseases and nitrogen deficiencies. These spatial networks demonstrated a quantum leap in accuracy compared to pixel-based shallow classifiers, because they evaluate a pixel's context within its neighbourhood rather than analysing its spectral signature in isolation.

A critical analysis of the current literature reveals several distinct research gaps that prevent the widespread adoption of UAV-deep learning pipelines in commercial agriculture:




FIGURE 1 Current Research Gaps in AI-Based Crop Growth Monitoring and Yield Prediction

First, the vast majority of existing deep learning models treat multi-temporal UAV datasets as isolated static events. Crops are dynamic biological systems; an anomaly observed in week eight is intrinsically linked to the developmental trajectory established in weeks two through seven. Models that rely purely on spatial CNNs miss these vital temporal dynamics. Second, there is a distinct lack of end-to-end integration between crop health diagnostic models and predictive yield models. Most frameworks address these as separate tasks, failing to realize that localized disease or nutrient stress early in the season has a non-linear compounding effect on final yield. Finally, many published models are overfit to small and well-localized active training datasets, making them impractical when deployed across varying crop varieties, soils, or illumination conditions. To directly overcome these limitations, this research defines a unified spatial-temporal deep learning network that models the entire crop lifecycle while maintaining strong generalizability and high prediction accuracy.

3. RESEARCH METHODOLOGY

This study was conducted in a methodical manner involving multiple levels of methodology to maximize data integrity, model robustness, and real-life applicability. The workflow includes five main components: data collection, image preprocessing, deep learning architecture design and model training, and evaluation.

3.1. DATA ACQUISITION AND FIELD EXPERIMENTAL DESIGN

Field experiments were undertaken over an entire agricultural-plant-to-harvest cultivating season on a large experimental farm with several diverse agro-ecological sectors. The crop used in this study, maize (*Zea mays* L.), was chosen for two reasons: 1) its structural complexity; and 2) as the most widely produced cereal grain worldwide. To obtain a more generalized deep learning model dataset, plots of the field were made with controlled treatments. Structured treatments varied nitrogen fertilizer application rates (low vs optimal vs excessive), water irrigation schedules (well-watered or not), and level of natural pest/disease exposure.

UAV flight missions were executed at critical phenological growth stages: seedling, V6 (six-leaf stage), V12, tasseling, silking, and physiological maturity. Flights were performed using a professional quadcopter UAV platform equipped with an integrated payload consisting of a high-resolution 20-megapixel RGB digital camera and a synchronized five-band multispectral sensor (capturing Blue, Green, Red, Red Edge, and Near-Infrared wavelengths). To minimize variations in solar illumination and shadow casting, all flights were executed under clear-sky conditions within two hours of local solar noon. Flights were kept at an altitude of 50 m AGS, which resulted in average ground sampling distance (GSD) values of 1.2 cm/pixel for RGB images and up to 4.5 cm/pixel for multispectral bands. Transverse and lateral overlaps were set to be 85% (for forward overlapping images) and 80%, respectively (photogrammetry), in order not to compromise the quality of reconstruction.

3.2. IMAGE PREPROCESSING AND PHOTOGRAMMETRIC RECONSTRUCTION

Each flight of data created raw aerial imagery that then passed through a rigorous preprocessing pipeline to create radiometrically calibrated, geometrically rectified orthomosaics from the initial raw digital numbers. The precise steps of the preprocessing workflow are detailed below:

- **Structure-from-Motion (SfM) Photogrammetry:** Raw images were imported into professional photogrammetry software where keypoints were automatically matched across overlapping frames to reconstruct the precise 3D geometry of the scene.
- **Ground Control Point (GCP) Alignment:** To ensure absolute geometric accuracy and perfect spatial alignment between multi-temporal flights, a set of ten highly visible, permanent Ground Control Points distributed across the field boundaries were surveyed using a differential Global Navigation Satellite System (GNSS) with Real-Time Kinematic (RTK) corrections, achieving sub-centimeter positioning accuracy.
- **Radiometric Calibration:** Raw multispectral digital numbers were converted into surface reflectance values utilizing downwelling light sensor data captured during the flight, alongside pre- and post-flight imagery of a calibrated radiometric calibration target with known reflectance values.
- **Orthomosaic Generation -** Using the calibrated images, RGB and multispectral ortho-mosaics were generated by naively projecting features onto the newly constructed DSMs for each observation window.
- **Tiling and Data Augmentation:** Large orthomosaics were tiled into 256×256 pixel patches that are applied homogeneously. We utilized data augmentation (random rotations, vertical/horizontal flips, and slight brightness changes) only on the training subsets to reduce overfitting and simulate a larger diversity of samples.

3.3. DEEP LEARNING ARCHITECTURE DESIGN

At the heart of this algorithm is the realization of a custom Hybrid Spatial-Temporal Deep Learning Network, specifically designed to extract spatial-spectral patterns from single data and monitor crop development through time. The architecture of the network is divided into two operational parts. The first module has an architecture that is modified from MS-CNN to take spatial inputs. A tile patch of RGB, single multispectral bands, and pre-computed vegetation index (i.e., NDVI and NDRE) layers are input into parallel convolutional streams. This design allows the network to capture independent fine structural details from high-resolution RGB imagery while simultaneously absorbing deep spectral signatures from multispectral layers. This optimization scheme is composed of all parallel feature maps generated by the proposed merge method using a synthesized feature layer with channel attention to weight spectral-spatial bands that are most diagnostic for each growth stage and healthy crop state.

The Second Module is a Recurrent Temporal network with Long Short-Term Memory (LSTM) blocks. For each phenological stage, the MS-CNN extracts spatial-spectral feature vectors organized chronologically and sequentially as inputs to LSTM layers. The LSTM architecture with internal memory gate approaches enables it to remember the growth velocity and stress progression of a crop over an entire season. The very last layer of the network has a dual-headed output configuration: a softmax classification head for obtaining a probability distribution over specific crop stresses (Healthy, Nitrogen Deficient, Water Stressed, or Diseased) and a parallel regression head as fully connected dense layers that outputs a continuous scalar value representing predicted metric tons per hectare with 95% confidence on complex networks.

4. RESULTS AND DISCUSSION

We investigated the efficacy of our proposed hybrid deep learning model by training and validating it against conventional machine learning classifiers as well as standard spatial-only CNN models for crop stress classification (yield level prediction). The deep learning model was developed using open-source software and trained on a high-performance computing workstation with multi-node NVIDIA tensor-core GPUs.

4.1. CROP STRESS CLASSIFICATION ANALYSIS

An independent validation dataset (not included in any of the four training data partitions) was prepared and used to rigorously evaluate model performance on classifying crop health conditions. Each crop tile was categorized into one of the four classes through a classification task: Healthy, Nitrogen-deficient, Water-stressed, or Diseased. Table 1 presents class-wise performance metrics obtained by the proposed hybrid architecture and therefore gives an overall evaluation of our networks.

TABLE 1 Detailed Class-Wise Classification Performance Metrics

Target Class	Precision (%)	Recall (%)	F1-Score (%)	Support (Samples)
Healthy	98.2	97.5	97.8	1,250
Nitrogen Deficient	95.4	96.1	95.7	980
Water Stressed	96.8	94.9	95.8	850
Diseased	94.7	96.3	95.5	720
Overall Average / Accuracy	96.4	96.3	96.3	3,800

As shown in Table 1, the statistical results indicate high sensitivity and specificity for predicting each of the health states with an overall classification accuracy rate of 96.4%. It is important to note that the model reached a very high precision within the Healthy class (98.2%), meaning only 1.8% of all best conditions were marked as positive by our classification system. In contrast, in reality they were negative, i.e., false positives when labelling optimal crop growing data points. Most importantly, the nitrogen deficient and water-stressed classes obtain an F1-score of 95.7% and 95.8%, respectively. It is notoriously difficult to separate nutrient shortages from moisture deficits, because both stressors appear phenotypically as chlorosis (yellowing of leaves). The success of the proposed network can be traced back to integrating Red Edge and Near-Infrared bands in the multi-stream convolutional layer, which hold essential information on cellular structural changes as well as early chlorophyll loss prior to other visible leaf regions observed by RGB imagery. With a maximum recall of 96.3% for the 'Diseased' class, the model exhibited high performance in localized pathogen detection, which is imperative to achieve target-specific automated pesticide deployment.

4.2. CROP YIELD PREDICTION AND COMPARATIVE ASSESSMENT

To validate the yield prediction capabilities of the regression head, the final outputs generated by the network at physiological maturity were compared against actual spatial yield data collected directly by automated grain yield monitors integrated into mechanical harvesters. The performance was quantified using standard statistical metrics: Coefficient of Determination (R^2), Root Mean Squared Error (RMSE) in tons per hectare (t/ha), and Mean Absolute Error (MAE).

To contextualize the performance of our proposed Hybrid MS-CNN + LSTM framework, three baseline models were trained on identical datasets: a standard Random Forest (RF) regressor utilizing hand-crafted VIs and textures, a Support Vector Machine (SVM), and a standard spatial-only CNN (ResNet-50) operating exclusively on late-season orthomosaics. Table 2 shows the comparative evaluation metrics of our proposed method.

TABLE 2 Comparative Yield Prediction Performance of Different Models

Model Architecture	Input Data Modality	R^2 Coefficient	RMSE (t/ha)	MAE (t/ha)
Random Forest (RF)	Hand-crafted VIs + GLCM Textures	0.71	1.42	1.12
Support Vector Machine (SVM)	Multispectral Pixel Values	0.68	1.55	1.24
Standard CNN (ResNet-50)	Single-date Late Season Orthomosaic	0.81	1.04	0.82
Proposed Hybrid Model	Multi-temporal Spatial-Spectral Fusion	0.89	0.62	0.45

It is clear from the comparative data shown in Table 2 that all other architectures evaluate poorly against the proposed Hybrid Model. The fewest errors were shown by the shallow machine learning models, SVM ($R^2 = 0.68$) and Random Forest ($R^2 = 0.71$). This underperformance highlights the deficiencies of manual feature engineering and conventional vegetation indices, which saturate during periods with high canopy density but do not account for more complex spatial interactions. The standard spatial-only CNN achieved a respectable R^2 of 0.81, demonstrating the inherent power of deep convolutional layers in capturing complex canopy textures and spatial layouts automatically. However, because the standard CNN processes only a single late-season orthomosaic, it remains oblivious to early-season stresses that permanently alter the crop's reproductive capacity.

The proposed Hybrid Network achieved the highest predictive accuracy, with an R^2 coefficient of 0.89, an RMSE of 0.62 t/ha , and an MAE of 0.45 t/ha . This improvement highlights the value of integrating temporal modelling via LSTM blocks. This network uses sequential growth data from a number of previous stages to estimate historical stresses, which helps determine their impact on the end development grain.




FIGURE 2 Actual vs. Predicted Crop Yield Comparison for the Proposed Machine Learning Model

This capability allows the model to differentiate between a field sector that is healthy but delayed in development and one suffering from severe, irreversible physiological degradation, providing accurate yield forecasts weeks before harvesting operations begin.

5. CONCLUSION

This work has successfully designed, implemented, and validated a smart agriculture framework based on ultra-high spatial resolution UAV imagery coupled with an original hybrid deep learning architecture. The proposed solution solves the main obstacles towards building operational agricultural remote sensing workflows by unifying multi-stream convolutional networks with temporal long short-term memory layers. Combining high-resolution spatial details with multispectral vegetation indices allowed our fully automated system to successfully characterize intra-field variability, partition highly correlated physiological stresses, and predict final crop yields with a consistently close match.

Experimental results confirmed that incorporation of spatial, spectral, and temporal data streams outperformed conventional shallow classifiers or single-date spatial networks alone. Achieving a crop stress classification accuracy of 96.4% and an R^2 yield prediction score of 0.89 demonstrates that this framework offers a highly reliable method for automated field diagnostics. Practically, this framework gives agronomists and large-scale farming enterprises the capacity to transition from historical reactive management to proactive, prescriptive precision farming. This shift enables variable-rate input applications, early localized disease control, and highly accurate supply chain planning with reliable pre-harvest yield predictions.

6. FUTURE SCOPE

Although the framework presented in this study provides state-of-the-art predictive accuracy and diagnostic precision, many exciting research directions remain for expanding its utility, scalability, and performance:

- **Fusion of multimodal remote sensing data:** Future work will focus on a fusion between high-resolution imagery identified from UAVs (Unmanned Aerial Vehicles) with continuous satellite streams (e.g., Sentinel-2), and ground-based Internet of Things (IoT) soil moisture/microclimate sensor networks to enable true multi-scale, continuous monitoring techniques across fields.
- **Optimization of Edge Computing for Real-Time Diagnostics:** Future research will focus on model compression techniques (weight quantization, network pruning, knowledge distillation) allowing complex deep learning networks to operate efficiently in situ as low-power edge computing hardware mounted directly onto the Autonomous UAV platform, enabling real-time decision making.
- **Cross-Crop and Multi-Geographic Validation:** Training and validating the architecture across various crop species (e.g., wheat versus rice versus specialized horticultural crops) and soil-climate geographic zones will be essential in making a truly global
- **Autonomous Variable-Rate Prescription Generative Pipelines:** Developing an automated software bridge that translates the output stress maps directly into standard shapefiles compatible with commercial tractor machinery will close the loop from diagnostic detection to autonomous, variable-rate field remediation.

REFERENCES

- [1] S. Ahirwar, R. Swarnkar, S. Bhukya, and G. Namwade, "Application of Drone in Agriculture," *International Journal of Current Microbiology and Applied Sciences*, vol. 8, no. 01, pp. 2500–2505, Jan. 2019, doi: 10.20546/ijcmas.2019.801.264.
- [2] M. Bhandari et al., "Assessing the Effect of Drought on Winter Wheat Growth Using Unmanned Aerial System (UAS)-Based Phenotyping," *Remote Sensing*, vol. 13, no. 6, p. 1144, Mar. 2021, doi: 10.3390/rs13061144.
- [3] J. G. A. Barbedo, "A Review on the Use of Unmanned Aerial Vehicles and Imaging Sensors for Monitoring and Assessing Plant Stresses," *Drones*, vol. 3, no. 2, p. 40, Apr. 2019, doi: 10.3390/drones3020040.
- [4] T. Chen and C. Guestrin, "XGBoost: a Scalable Tree Boosting System," *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining - KDD '16*, vol. 1, no. 1, pp. 785–794, Aug. 2016, doi: 10.1145/2939672.2939785.
- [5] Kamilaris and F. X. Prenafeta-Boldú, "Deep learning in agriculture: A survey," *Computers and Electronics in Agriculture*, vol. 147, pp. 70–90, Apr. 2018, doi: 10.1016/j.compag.2018.02.016.
- [6] T. Kattenborn, J. Leitloff, F. Schiefer, and S. Hinz, "Review on Convolutional Neural Networks (CNN) in vegetation remote sensing," *ISPRS Journal of Photogrammetry and Remote Sensing*, vol. 173, pp. 24–49, Mar. 2021, doi: 10.1016/j.isprsjprs.2020.12.010.
- [7] Khan, A. D. Vibhute, S. Mali, and C. H. Patil, "A systematic review on hyperspectral imaging technology with a machine and deep learning methodology for agricultural applications," *Ecological Informatics*, vol. 69, p. 101678, July 2022, doi: 10.1016/j.ecoinf.2022.101678.
- [8] N. Kussul, M. Lavreniuk, S. Skakun, and A. Shelestov, "Deep Learning Classification of Land Cover and Crop Types Using Remote Sensing Data," *IEEE Geoscience and Remote Sensing Letters*, vol. 14, no. 5, pp. 778–782, May 2017, doi: 10.1109/lgrs.2017.2681128.
- [9] V. Singh, N. Sharma, and S. Singh, "A review of imaging techniques for plant disease detection," *Artificial Intelligence in Agriculture*, vol. 4, Oct. 2020, doi: 10.1016/j.aiaa.2020.10.002.
- [10] M. Maimaitijiang, V. Sagan, P. Sidike, S. Hartling, F. Esposito, and F. B. Fritschi, "Soybean yield prediction from UAV using multimodal data fusion and deep learning," *Remote Sensing of Environment*, vol. 237, p. 111599, Feb. 2020, doi: 10.1016/j.rse.2019.111599.

- [11] P. Radoglou-Grammatikis, P. Sarigiannidis, T. Lagkas, and I. Moscholios, "A Compilation of UAV Applications for Precision Agriculture," *Computer Networks*, vol. 172, p. 107148, Feb. 2020, doi: 10.1016/j.comnet.2020.107148.
- [12] V. Sagan et al., "UAV-Based High Resolution Thermal Imaging for Vegetation Monitoring, and Plant Phenotyping Using ICI 8640 P, FLIR Vue Pro R 640, and thermoMap Cameras," *Remote Sensing*, vol. 11, no. 3, p. 330, Feb. 2019, doi: 10.3390/rs11030330.
- [13] D. C. Tsouros, S. Bibi, and P. G. Sarigiannidis, "A Review on UAV-Based Applications for Precision Agriculture," *Information*, vol. 10, no. 11, p. 349, Nov. 2019, doi: 10.3390/info10110349.
- [14] M. Weiss, F. Jacob, and G. Duveiller, "Remote sensing for agricultural applications: A meta-review," *Remote Sensing of Environment*, vol. 236, p. 111402, Jan. 2020, doi: 10.1016/j.rse.2019.111402.
- [15] P. Muruganantham, S. Wibowo, S. Grandhi, N. H. Samrat, and N. Islam, "A Systematic Literature Review on Crop Yield Prediction with Deep Learning and Remote Sensing," *Remote Sensing*, vol. 14, no. 9, p. 1990, Apr. 2022, doi: 10.3390/rs14091990.